



AMERICAN PSYCHOLOGICAL ASSOCIATION

Eating Disorders



In a society that continues to prize thinness even as Americans become heavier than ever before, almost everyone worries about their weight at least occasionally. People with eating disorders take such concerns to extremes, developing abnormal eating habits that threaten their well-being and even their lives. This question-and-answer fact sheet explains how psychotherapy can help people recover from these dangerous disorders.

What are the major kinds of eating disorders?

There are three major types of eating disorders.

People with **anorexia nervosa** have a distorted body image that causes them to see themselves as overweight even when they're dangerously thin. Often refusing to eat, exercising compulsively, and developing unusual habits such as refusing to eat in front of others, they lose large amounts of weight and may even starve to death.

Individuals with **bulimia nervosa** eat excessive quantities, then purge their bodies of the food and calories they fear by using laxatives, enemas, or diuretics; vomiting; or exercising. Often acting in secrecy, they feel disgusted and ashamed as they binge, yet relieved of tension and negative emotions once their stomachs are empty again.

Like people with bulimia, those with **binge eating disorder** experience frequent episodes of out-of-control eating. The difference is that binge eaters don't purge their bodies of excess calories.

Another category of eating disorders is "eating disorders not otherwise specified," in which individuals have eating-related problems but don't meet the official criteria for anorexia, bulimia or binge eating.

It's important to prevent problematic behaviors from evolving into full-fledged eating disorders. Anorexia and bulimia, for example, usually are preceded by very strict dieting and weight loss. Binge eating disorder can begin with occasional bingeing. Whenever eating behaviors start having a destructive impact on someone's functioning or self-image, it's time to see a highly trained mental health professional, such as a licensed psychologist experienced in treating people with eating disorders.

Who suffers from eating disorders?

According to the National Institute of Mental Health (<http://www.nimh.nih.gov/health/publications/eating-disorders/how-are-males-affected.shtml>), eating disorders primarily affect girls and women.¹ (#national) But eating disorders aren't just a problem for the teenage women so often depicted in the media. Men and boys can also be vulnerable. About a quarter of preadolescent cases of anorexia occur in boys, for example. And binge eating disorder strikes males and females about equally. People sometimes have eating disorders without their families or friends ever suspecting that they have a problem. Aware that their behavior is abnormal, people with eating disorders may withdraw from social contact, hide their behavior, and deny that their eating patterns are problematic. Making an accurate diagnosis requires the involvement of a licensed psychologist or other appropriate mental health expert.

What causes eating disorders?

Certain psychological factors and personality traits may predispose people to developing eating disorders. Many people with eating disorders suffer from low self-esteem, feelings of helplessness, and intense dissatisfaction with the way they look.

Specific traits are linked to each of the disorders. People with anorexia tend to be perfectionistic, for instance, while people with bulimia are often impulsive. Physical factors such as genetics also may play a role in putting people at risk.

A wide range of situations can precipitate eating disorders in susceptible individuals. Family members or friends may repeatedly tease people about their bodies. Individuals may be participating in gymnastics or other sports that emphasize low weight or a certain body image. Negative emotions or traumas such as rape, abuse, or the death of a loved one can also trigger disorders. Even a happy event, such as giving birth, can lead to disorders because of the stressful impact of the event on an individual's new role and body image.

Once people start engaging in abnormal eating behaviors, the problem can perpetuate itself. Bingeing can set a vicious cycle in motion, for instance, as individuals purge to rid themselves of excess calories and psychic pain, then binge again to escape problems in their day-to-day lives.

Why is it important to seek treatment for these disorders?

Research indicates that eating disorders very often go untreated. In one study (<http://download.journals.elsevierhealth.com/pdfs/journals/0890-8567/PIIS0890856710007835.pdf>) (PDF, 382KB), for example, less than 13 percent of adolescents with eating disorders received treatment.² (#merikangas)

But leaving eating disorders untreated can have serious consequences. Research (<http://linkinghub.elsevier.com/retrieve/pii/S1056499308000631>) has found that individuals with anorexia have a mortality rate 18 times higher than peers who don't have eating disorders, for example.³ (#steinhausen)

Eating disorders can devastate the body. Physical problems associated with anorexia, for instance, include anemia, constipation, osteoporosis, even damage to the heart and brain. Bulimia can result in a sore throat, worn-away tooth enamel, acid reflux, and heart attacks. People with binge eating disorder may develop high blood pressure, cardiovascular disease, diabetes, and other problems associated with obesity.

Eating disorders are also associated with other mental disorders like depression. Researchers don't yet know whether eating disorders are symptoms of such problems or whether the problems develop because of the isolation, stigma, and physiological changes wrought by the eating disorders themselves. What is clear from the research (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1892232/pdf/nihms19542.pdf>) (PDF, 399KB) is that people with eating disorders suffer higher rates of other mental disorders - including depression, anxiety disorders, and substance abuse - than other people.⁴ (#hudson)

How can a psychologist help someone recover?

Psychologists play a vital role in the successful treatment of eating disorders and are integral members of the multidisciplinary team that may be required to provide patient care. As part of this treatment, a physician may be called on to rule out medical illnesses and determine that the patient is not in immediate physical danger. A nutritionist may be asked to help assess and improve nutritional intake.

Once the psychologist has identified important issues that need attention and developed a treatment plan, he or she helps the patient replace destructive thoughts and behaviors with more positive ones. A psychologist and patient might work together to focus on health rather than weight, for example. Or a patient might keep a food diary as a way of becoming more aware of the types of situations that trigger bingeing.

Simply changing patients' thoughts and behaviors is not enough, however. To ensure lasting improvement, patients and psychologists must work together to explore the psychological issues underlying the eating disorder. Psychotherapy may need to focus on improving patients' personal relationships. And it may involve helping patients get beyond an event or situation that triggered the disorder in the first place. Group therapy also may be helpful.

Some patients, especially those with bulimia, may benefit from medication. It's important to remember, however, that medication should be used in combination with psycho-therapy, not as a replacement for it. Patients who are advised to take medication should be aware of possible side effects and the need for close supervision by a physician.

Does treatment really work?

Yes. Most cases of eating disorder can be treated successfully by appropriately trained health and mental health care professionals. But treatments do not work instantly. For many patients, treatment may need to be long-term.

Incorporating family or marital therapy into patient care may help prevent relapses by resolving interpersonal issues related to the eating disorder. Therapists can guide family members in understanding the patient's disorder and learning new techniques for coping with problems. Support groups can also help.

Remember: The sooner treatment starts, the better. The longer abnormal eating patterns continue, the more deeply ingrained they become and the more difficult they are to treat.

Eating disorders can severely impair people's functioning and health. But the prospects for long-term recovery are good for most people who seek help from appropriate professionals. Qualified therapists, such as licensed psychologists with experience in this area, can help those who suffer from eating disorders regain control of their eating behaviors and their lives.

Thanks to Kelly D. Brownell, PhD; Kathy J. Hotelling, PhD; Michael R. Lowe, PhD; and Gina E. Rayfield, PhD, who assisted with this article.

Revised October 2011

Article Sources

¹ National Institute of Mental Health. (2007). "Eating disorders."

² Merikangas, K.R., He, J., Burstein, M., Senden, J., Avenevoli, S., Case, B., Georgiades, K., et al. (2011). "Service utilization for lifetime mental disorders in U.S. adolescents: Results of the National Comorbidity Survey-Adolescent Supplement (NCS-A)." *Journal of the American Academy of Child & Adolescent Psychiatry*, 50 (1): 32-45.

³ Steinhausen, H.C. (2009). "Outcomes of eating disorders." *Child and Adolescent Psychiatric Clinics of North America*, 18 (1): 225-242.

⁴ Hudson, J.I., Hiripi, E., Pope, H.G., & Kessler, R.C. (2007). "The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication." *Biological Psychiatry*, 61 (3): 348-358.

The full text of articles from APA Help Center may be reproduced and distributed for noncommercial purposes with credit given to the American Psychological Association. Any electronic reproductions must link to the original article on the APA Help Center. Any exceptions to this, including excerpting, paraphrasing or reproduction in a commercial work, must be presented in writing to the APA.

(mailto:permissions@apa.org) Images from the APA Help Center may not be reproduced.

Find this article at:

<http://www.apa.org/helpcenter/eating.aspx>